

AniMove 2024, June 17th to 28th


Home range meta-analyses

Using the 'ctmm' R package

Inês Silva, Chris Fleming

✉ i.simoes-silva@hzdr.de

Analyses of ecological data should always account for the **uncertainty** in the process(es) that generated the data.



Accurately estimating **area requirements** is of utmost importance for conservation, from the **individual** to the **population level**.




Analyses of ecological data should always account for the **uncertainty** in the process(es) that generated the data.



Accurately estimating **area requirements** is of utmost importance for conservation, from the **individual** to the **population level**.



We want to quantify the effect of covariates, such as **species or taxa**, **sex**, **body size**, **age**, **movement characteristics**, **conspecific density**, **habitat**, **anthropogenic impact**, etc....



Analyses of ecological data should always account for the **uncertainty** in the process(es) that generated the data.



Accurately estimating **area requirements** is of utmost importance for conservation, from the **individual** to the **population level**.




We want to quantify the effect of covariates, such as **species or taxa**, **sex**, **body size**, **age**, **movement characteristics**, **conspecific density**, **habitat**, **anthropogenic impact**, etc....



... even if we are comparing different populations with **different movement behaviors** or **sampling schedules**.

NON-HIERARCHICAL MODELS


How does data inform parameters?



 Adapted from Midway (2008)

NON-HIERARCHICAL MODELS


How does data inform parameters?



Adapted from Midway (2008)

HIERARCHICAL MODELS


How does data inform parameters?



 Adapted from Midway (2008)


HIERARCHICAL MODELS

How does data inform parameters?




HIERARCHICAL MODELS

How does data inform parameters?




Trajectory



Data

χ^2 inverse-Gaussian meta-analysis


Individual



↓ $p(\theta_i | \Theta)$


↓ $p(\mathbf{r}_i | \theta_i)$

Trajectory






Population




$$\downarrow p(\theta_i | \Theta)$$


Individual




Tracking data



Home range analysis



Population analysis








LOWLAND TAPIR TAPIRUS TERRESTRIS



Tapirs have HR crossing times (τ_p) of 0.72 days, ranging from 0.05 to 12.8 days.





This framework facilitates population-level inference with as few as **2-3 observed home range crossings** (τ_p) and similarly small **number of individuals** (m).

